

**TRANSMITTAL**

To: Bureau of Design and Environment  
Attention: Tom Brooks  
From: Illinois Natural History Survey  
Regarding: Mitigation Monitoring Project

**Project Information**

Requesting Agency: IDOT Springfield  
County: Macon  
Route: FAP 322 (U.S. 51)  
Section: 58-20-1 & 58-20-1 (HB, HB-1, HB-2) BR & 48R-1  
At: Approximately 1 mi. south of Elwin, IL. Within the south infield at the U.S. 51-Riley Road Interchange.  
Quadrangle: Decatur, IL

**Survey Conducted By:** Paul Marcum and Jeff Matthews (vegetation and hydrology)  
Jessica Kurylo (soils and hydrology)  
Brad Zercher (GPS, GIS and hydrology)  
Eric Plankell and Geoff Pociask (ISGS; hydrology)  
Illinois Natural History Survey  
Division of Ecology and Conservation Science  
Wildlife and Plant Ecology Section  
1816 S. Oak Street  
Champaign, IL 61820  
(217) 333-8459 (Marcum), 244-0692 (Kurylo)

**Date Conducted:** 5 June and 29 August 2006

**Project Summary:**

Fifth and final year field monitoring of a wetland mitigation site along FAP 322 (U.S. 51) was conducted on 5 June and 29 August 2006. This site is located approximately 1 mile south of Elwin, IL within the south infield at the U.S. 51 - Riley Road interchange. The wetland is a created depression within this infield. An emergent wetland community (wet prairie/marsh) is planned for this site. The dominant species in the created marsh community is *Typha angustifolia*. In 2006, approximately 2.01 ha (4.96 ac) of the 4.7 ha (11.6 ac) site satisfied all three wetland criteria. This report documents the final year of monitoring for this mitigation area. Summary information regarding the wetland delineation sites is presented in the project report. The wetland determination forms are contained in Appendix 2 and photos of the wetland creation site are included in Appendix 3.

Signed: \_\_\_\_\_  
Dr. Allen E. Plocher  
INHS/IDOT Project Coordinator

Signed: \_\_\_\_\_  
Dr. Edward J. Heske  
INHS/IDOT Project Principal Investigator

Date: \_\_\_\_\_

Date: \_\_\_\_\_

**Fifth Year Wetland Mitigation Site Monitoring for FAP 322  
(U.S. 51), Macon County, Illinois – 2006**

Paul Marcum, Jessica Kurylo, Jeff Matthews, and Brad Zercher  
Illinois Natural History Survey  
Division of Ecology and Conservation Science  
1816 S. Oak Street  
Champaign, Illinois 61820  
(217) 333-8459 (Marcum)

**Introduction**

Wetland compensation activity has been initiated along U.S. 51 in Macon County, Illinois. The legal location of the site is NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E. (Decatur, IL Quad). Prior to road construction this area had been farmed for over 75 years and cover consisted of corn and beans (Coopriider and Ketzner 2000). The pre-settlement environment consisted of wet and mesic prairie. Presently, the site is surrounded by road embankments and is mostly a depression within the infield of the U.S. 51 – Riley Road interchange. The mitigation site assessment and wetland compensation plan for this area suggested that an emergent wetland plant community (i.e. wet prairie, wet meadow, or marsh) would be the most likely development for this site (Coopriider and Ketzner 2000; IDOT 1996). Approximately 150 m (500 ft) west of the wetland mitigation site is the Elwin *Camassia* site (Brooks 1999; IDOT 1996). This site is a mesic to dry-mesic prairie Illinois Natural Areas Inventory site that contains a population of the state endangered wild hyacinth, *Camassia angusta* (Engelmann & A. Gray) Blankinship (Herkert and Ebinger 2002).

According to Coopriider and Ketzner (2000), 0.79 ha (1.95 ac) of this infield was already wetland in 1999. Since then, the site has been shallowly excavated, mostly at the northern end of the infield, creating more low depressional ground to support a wetland. An emergent wetland community (wet prairie/marsh) is planned for this site. Construction of this wetland mitigation site was completed in June 2001. Beginning in 2002, Illinois Natural History Survey (INHS) personnel have monitored this area for five years as requested by the Illinois Department of Transportation (Brooks 2001). The Illinois State Geological Survey (ISGS) has also monitored the hydrology of this site. Project goals, objectives, and performance criteria are included in this report, as are monitoring methods, monitoring results, summary information and recommendations.

**Project Goals, Objectives, and Performance Criteria**

Proposed goals and objectives for this wetland mitigation project are based on information contained in the original wetland compensation plan for this site (IDOT 1996). Performance criteria are based on those specified in the *Corps of Engineers Wetland Delineation Manual* (Environmental Laboratory 1987) and *Guidelines for Developing Mitigation Proposals* (USACOE 1993). Each goal should be attained by the end of the five-year monitoring period. Project goals, objectives and performance criteria are listed below.

**Project Goal #1:** At the end of the five-year monitoring period the created wetland community should be a jurisdictional wetland as defined by current federal standards.

**Objective:** The created wetland should compensate for the loss of wetland.

**Performance Criteria:** The entire created wetland should satisfy the three criteria of the federal wetland definition: dominant hydrophytic vegetation, hydric soils, and wetland hydrology.

- A. Predominance of Hydrophytic Vegetation – More than 50% of the dominant plant species must be hydrophytic.
- B. Presence of Hydric Soils – Hydric soil characteristics should be present, or conditions favorable for hydric soil formation should persist at this site.
- C. Presence of Wetland Hydrology – The compensation area must be either permanently or periodically inundated at average depths less than 2 m (6.6 ft) or have soils that are saturated to the surface for at least 12.5% of the growing season.\*

**Project Goal #2:** A native, non-weedy, emergent wetland community will be created.

**Objective:** Planting the area with high quality native emergent vegetation should reduce the pressures from successional, non-native, weedy species.

**Performance Criteria:** At least 50% of the plant species present should be non-weedy, native, perennial species. Furthermore, none of the dominant plant species may be non-native, cattails, or reed canary grass.

## Methods

Monitoring of this wetland mitigation site began in 2002 and has continued for the standard five-year monitoring period. Illinois Natural History Survey (INHS) personnel have monitored the biological parameters and Illinois State Geological Survey (ISGS) personnel have monitored hydrology. Herbaceous vegetation has been assessed annually using standard sampling techniques (Cox 1985). Transects were established through the created wetland with a minimum of 30 1m<sup>2</sup> quadrats sampled annually. Results and status of the created wetland site have been submitted to the Illinois Department of Transportation (IDOT) in yearly monitoring reports. The likelihood of meeting the proposed goals and performance criteria has been addressed. If, at any time during the monitoring period, it appeared that the goals/performance criteria would not be met at the end of the five-year monitoring period, written management recommendations were made to IDOT in an effort to correct any problems.

## Floristic Quality Index

A complete list of all plant species found in the project area was recorded annually and the Floristic Quality Index (FQI) was calculated (Swink and Wilhelm 1979 and 1994; Taft *et al.*

---

\* In some cases wetland hydrology can be met when a site is inundated or saturated for 5% to 12.5% of the growing season (Environmental Laboratory 1987).

1997). The FQI provides a measure of the floristic integrity or level of disturbance of a site. Each native plant species is assigned a rating between 0 and 10 (the Coefficient of Conservatism) that is a subjective indicator of the likelihood that a plant may be found on an undisturbed site in a natural plant community. A plant species that has a low Coefficient of Conservatism (C) is common and is likely to tolerate disturbed conditions; a species with a high C is relatively rare and is likely to require specific, undisturbed habitats. Species not identified to species level are not rated and are not included in the calculations.

To calculate the FQI, first compute the mean C value (also known as mean rated quality),  $mCv = \sum C/N$ , where  $\sum C$  represents the sum of the numerical ratings (C) for all species recorded for a site, and N represents the number of plants on the site. The C value for each species is shown in the species list for the site. Species that are not native to Illinois (indicated by \* in the species list for each site) are not included in the calculations. The FQI for each site is determined by dividing the  $\sum C$  value by the square root of N [ $\sum C/(\sqrt{N})$ ]. An Index score below 10 suggests a site of low natural quality; below 5, a highly disturbed site. An FQI value of 20 or more ( $mCv > 3$ ) suggests that a site has evidence of native character and may be considered an environmental asset.

**Project Goal #1** At the end of the five year monitoring period the created wetland community should be a jurisdictional wetland as defined by current federal standards.

Wetland delineation has been completed yearly for all wetland community types at this compensation site. In addition, permanent photo stations have been established in the wetland restoration/creation area and photos have been taken annually in order to help monitor changes in the vegetation. Photos are included in Appendix 3 of the report.

A. **Predominance of Hydrophytic Vegetation** – The method for determining dominant hydrophytic vegetation is described in Environmental Laboratory (1987) and Federal Interagency Committee for Wetland Delineation (1989). This method is based on aerial coverage estimates for individual plant species. Each of the dominant plant species is assigned a wetland indicator status rating (Reed 1988). Any plant rated facultative or wetter (i.e., FAC, FAC+, FACW-, FACW, FACW+ and OBL) is considered hydrophytic. A predominance of hydrophytic vegetation in the wetland plant community exists if greater than 50% of the dominant species present are hydrophytic.

Dominant hydrophytic vegetation has been determined each year based on the results of systematic plant sampling. Cover of all species in each plot is assigned a cover class according to Daubenmire (1959) (Table 1). Frequency (proportion of quadrats in which a species occurred) and average cover (calculated using midpoints for each cover class) is used to compute relative frequency (frequency of a species relative to total observations) and relative cover (cover relative to total observed cover), respectively. These two relative values are averaged to determine the importance value for each species sampled. Importance values will be used to determine dominant species. “Dominant species are the most abundant plant species (when ranked in descending order of abundance and cumulatively totaled) that immediately exceed 50% of the total dominance measure for the stratum, plus any additional species comprising 20% or more of the total dominance measure for the stratum” (FICWD 1989; Tiner 1999).

**Table 1. Cover classes used in vegetation sampling**

Cover Class	Range of Cover (%)	Midpoint of Range (%)
1	0-5	2.5
2	5-25	15.0
3	25-50	37.5
4	50-75	62.5
5	75-95	85.0
6	95-100	97.5
(Daubenmire 1959)		

B. Presence of Hydric Soils – Soils have been examined and described annually. A soil core collected from the same general area of the mitigation site has been examined for the presence of redoximorphic features. A detailed profile description of the soil using Munsell color charts to record soil colors is included. Soil texture and structure is also recorded. Hydric soils may develop slowly and characteristics may not be apparent during the first several years after project construction. In the absence of hydric soil indicators at that time, hydrologic data could be used as corroborative evidence that conditions favorable for hydric soil formation are present at the site.

C. Presence of Wetland Hydrology – The ISGS has been tasked to monitor hydrology at the proposed wetland site. To date they have installed two surface-water monitoring stations (RDS1 and RDS2), a rain guage, two surface-water staff gauges (A and B), and eleven monitoring wells (1S-8S and 10S-12S) (Watson and Sabatini 2002; Watson and Sabatini 2003; Plankell and Pociask 2004; Plankell and Pociask 2005; Plankell and Pociask 2006). ISGS personnel measure water levels monthly. In addition, INHS scientists survey the site annually for field indicators of wetland hydrology.

**Project Goal #2** A native, non-weedy, emergent wetland community will be created.

**Performance Criteria:** At least 50% of the plant species present should be non-weedy, native, perennial species. Furthermore, none of the dominant plant species may be non-native, cattails, or reed canary grass.

A complete species list has been compiled each year and species are recorded as native or non-native, weedy or non-weedy, and as a perennial, biennial, or an annual. Nativity of plants is determined by consulting Mohlenbrock (1986; 2002) and Taft *et al.* (1997). Weedy species, for the purposes of this report, are defined as all non-native species and any native species assigned a Coefficient of Conservatism of 0 or 1 (Taft *et al.* 1997). Species given a C value of 0-1 correspond to Grime’s ruderal species (Grime 1974; Grime *et al.* 1988) or species which are adapted to frequent or severe disturbances (Taft *et al.* 1997). Gleason and Cronquist (1991), Taft *et al.* (1997), and USDA, NRCS (2002) were the primary sources used to determine whether a species is perennial, biennial, or annual.

## Results

**Floristic Quality Index:** The Floristic Quality Index was calculated for this site using native species only. The wetland creation/restoration site had a FQI of 16.7 and a mean C value of 2.4. These values are indicative of fair natural quality. The upland buffer around the wetland site had an FQI of 16.8 and a mean C value of 2.5. Likewise, these values are characteristic of fair natural quality. In 2006, a total of 49 native species were found in the wetland community with a few species indicative of higher natural quality being present. Species present with a C value of 5 or greater: *Aster puniceus* (7), *Iris shrevei* (5), *Lobelia cardinalis* (6), *Potamogeton pectinatus* (5), *Pycnanthemum virginianum* (5), *Solidago ridellii* (7), and *Sparganium eurycarpum* (5). Summary information for the created wetland site is given in Table 2.

**Table 2. Summary table for FAP 322 wetland monitoring site, 2002 to 2006.**

	2002	2003	2004	2005	2006
Total Species Richness	40	51	51	58	60
Native Species Richness	24	38	41	43	49
% Native	60%	75%	80%	74%	82%
% Non-weedy	23%	43%	51%	47%	48%
% Perennial	35%	47%	71%	60%	68%
% Native, Non-weedy, and Perennial	18%	29%	45%	38%	43%
Mean Conservatism	1.6	2.2	2.5	2.5	2.4
Floristic Quality Index (FQI)	7.8	13.6	16.0	16.6	16.7
% Wetland Species (FAC to OBL)	60%	78%	82%	78%	82%

**Project Goal #1** At the end of the five year monitoring period the created wetland community should be a jurisdictional wetland as defined by current federal standards.

A. Predominance of Hydrophytic Vegetation – The performance criterion requires that greater than 50% of the dominant plant species be hydrophytic. Results for 2006 indicate that the dominant herbaceous species is *Typha angustifolia* (OBL) (Table 3). Greater than 50% (100%) of the dominant plant species are hydrophytic; therefore, this site meets the criterion for predominance of hydrophytic vegetation.

**Table 3. FAP 322 (U.S. 51) Wetland mitigation site vegetation sampling data including frequency, cover, and importance value for all species sampled in 2006 (dominant species is in bold).**

Species	Indicator	Frequency	Relative Frequency	Average Cover	Relative Cover	Importance Value
<b><i>Typha angustifolia</i></b>	<b>OBL</b>	<b>0.8750</b>	<b>46.6667</b>	<b>23.6875</b>	<b>60.1587</b>	<b>53.4127</b>
<i>Aster pilosus</i>	FACU-	0.1250	6.6667	3.8542	9.7884	8.2275
<i>Bidens cernua</i>	OBL	0.0833	4.4444	4.1667	10.5820	7.5132
<i>Aster puniceus</i>	OBL	0.0833	4.4444	3.6667	9.3122	6.8783
<i>Scirpus taebertmontanii</i>	OBL	0.1667	8.8889	1.0000	2.5397	5.7143
<i>Solidago canadensis</i>	FACU	0.1250	6.6667	1.3750	3.4921	5.0794
<i>Rumex crispus</i>	FAC+	0.1250	6.6667	0.3750	0.9524	3.8095
<i>Eleocharis erythropoda</i>	OBL	0.0833	4.4444	0.2500	0.6349	2.5397
<i>Asclepias incarnata</i>	OBL	0.0417	2.2222	0.6250	1.5873	1.9048
<i>Hordeum jubatum</i>	FAC+	0.0417	2.2222	0.1250	0.3175	1.2698
<i>Juncus torreyi</i>	FACW	0.0417	2.2222	0.1250	0.3175	1.2698
<i>Leersia oryzoides</i>	OBL	0.0417	2.2222	0.1250	0.3175	1.2698
<i>Ambrosia artemisiifolia</i>	FACU	0.0417	2.2222	0.0000	0.0000	1.1111
bare ground and litter		1.8750	100.0000	39.3750	100.0000	100.0000
				60.6250		

**Dominant species is in bold**

B. Presence of Hydric Soils – The performance criterion requires that hydric soil characteristics be present, or conditions favorable for hydric soil formation should persist. The soils on this site in 1999 were mapped as Drummer silty clay loam and Flanagan silt loam (Coopridier 1999). Since then, some of the non-hydric areas of this site have been excavated and altered to lower the surface. This is most evident in the northern end of the site where the substratum of the somewhat poorly drained soil (Flanagan) was exposed leading to the appearance of weak redox features at the surface. Since creation of this site, these soils have started to develop characteristics of their own based on their new pedogenic conditions, therefore expanding the hydric soil boundaries outward. As long as the hydrology is present, this area will remain hydric. At the time of the field visit most of the site was inundated. A typical pedon for this wetter area of the site is described below.

**Table 4. Description of the wetland soils at the FAP 322 wetland monitoring site.**

Depth(in)	Matrix Color	Concentrations	Depletions	Texture	Structure
0-1	10YR 2/1			Silty Clay Loam	Granular
1-15	2.5Y 4/2	10YR 5/8	10YR 4/1	Silty Clay Loam	Subangular Blocky
15-20	2.5Y 5/2 & 10YR 5/8	N 2.5/0		Silty Clay Loam	Subangular Blocky

In the northernmost and southernmost areas of the interchange and along US 51 the soils have little to no chance of becoming hydric due to their landscape position. A soil sample from the upland in the southern part of the project area is also described below.

**Table 5. Description of the non-wetland soils at the FAP 322 wetland monitoring site.**

Depth(in)	Matrix Color	Concentrations	Depletions	Texture	Structure
0-6	10YR 2/1	7.5 YR 5/8		Silty Clay Loam	Angular Blocky to Platy
6-13	10YR 6/6	10 YR 4/6		Silty Clay Loam	Weak Sub-Blocky to Massive

C. Presence of Wetland Hydrology – The performance criterion requires that the compensation area must be either permanently or periodically inundated at average depths less than 2m (6.6 ft) or have soils that are saturated to the surface for at least 12.5% of the growing season\*. The ISGS initiated water level monitoring at this site in September 2000. Their findings for 2006 indicate that 2.3 ha (5.7 ac) out of a total site area of approximately 4.7 ha (11.6 ac) satisfied the wetland hydrology criterion for greater than 12.5% of the growing season (Plankell and Pociask 2006; Figure 1). The area of satisfactory wetland hydrology has increased since 2005 [2.1 ha (5.2 ac)] but is somewhat less than 2002 [3.0 ha (7.4 ac)], 2003 [2.6 ha (6.3 ac)], and 2004 [2.4 ha (5.9 ac)] (Watson and Sabatini 2002; Watson and Sabatini 2003; Plankell and Pociask 2004; Plankell and Pociask 2005).

During visits to the site, the following indicators of wetland hydrology were present: areas of inundation, algal mats, mud cracks, drift, blackened leaves, and some areas of surface or near surface saturation.

**Project Goal #2:** A native, non-weedy, emergent wetland community will be created.

Many weedy native and non-native species were present during the first few years of sampling. This has improved somewhat, over the course of the past five years, as perennial species have become established. In 2006, 49 out of 60 (82%) plant species found at this site were native, 29 (48%) species are considered non-weedy, and 41 (68%) were perennials. However, when combined only 43% (26/60) of the species found at this wetland site were native, non-weedy, and perennial. Although the vegetation composition present at this site continues to improve, project goal #2 was not satisfied in 2006. Furthermore, the site continues to be dominated by *Typha angustifolia*, the non-native, narrow-leaf cattail.

---


\* In some cases wetland hydrology can be met when a site is inundated or saturated for 5% to 12.5% of the growing season (Environmental Laboratory 1987).

## Decatur, U.S. Route 51 Wetland Compensation Site (FAP 322)

### Estimated Areal Extent of 2006 Wetland Hydrology

Based on data collected between September 1, 2005 and September 1, 2006

Map based on USGS digital orthophotograph, Decatur SW quarter quadrangle  
produced from 4/14/98 aerial photography (ISGS 2000)



**Figure 1. 2006 aerial extent of wetland hydrology for FAP 322 wetland monitoring site (prepared by ISGS, Plankell and Pociask 2006)**

## Summary and Recommendations

**Floristic Quality Index** – The FQI has continued to increase since the first year of monitoring (Marcum *et al.* 2002). The FQI score for this wetland mitigation site has increased from a low of 7.8 in 2002 to a high of 16.7 in 2006 (Marcum *et al.* 2002; Marcum and Kurylo 2003; Marcum *et al.* 2004, Marcum *et al.* 2005). Likewise, the mean C value has increased from 1.6 in 2002 to 2.4 in 2006. These current values are indicative of fair natural quality. Similarly, the proportion of perennial species at this site has increased over time, resulting in a more stable plant community. The wetland compensation plan (IDOT 1996) states that the created wetland should be aesthetically appealing from the road. Planting of additional high quality species to the perimeter of the wetland and to parts of the upland buffer have helped somewhat in this regard.

**Project Goal # 1** – At the end of the five-year monitoring period the created wetland community should be a jurisdictional wetland as defined by current federal standards. The performance criterion requires that greater than 50% of the dominant plant species be hydrophytic, that hydric soil characteristics be present, or conditions favorable for hydric soil formation should persist, and that the compensation area must be either permanently or periodically inundated at average depths less than 2m (6.6 ft) or have soils that are saturated to the surface for at least 12.5% of the growing season\*. In 2006, this created/restored wetland site exhibited dominant hydrophytic vegetation, hydric soils, and wetland hydrology.

The area satisfying wetland hydrology was found to be somewhat more than last year but somewhat less than the first three years (Plankell and Pociask 2006; Plankell and Pociask 2005; Plankell and Pociask 2004; Watson and Sabatini 2003; Watson and Sabatini 2002). In 2002, the area of wetland hydrology was at a high point of 3.0 ha (7.4 ac), while in 2006, 2.3 ha (5.7 ac) out of a total site area of approximately 4.7 ha (11.6 ac) satisfied the wetland hydrology criterion for greater than 12.5% of the growing season (Watson and Sabatini 2002; Plankell and Pociask 2006; Figure 1).

INHS data from vegetation sampling, soil mapping, and general observations of wetland hydrology determine the aerial extent of the created wetland to be approximately 2.01 ha (4.96 ac) (Appendix 1, Figure 2). Furthermore, one area at the south end of the wetland [~ 150 ft by 150 ft or 0.21 ha (0.52 ac)] currently exhibits hydric soil development and satisfies wetland hydrology. This area, however, still has yet to develop hydrophytic vegetation because of the dominance of meadow fescue, *Festuca pratensis*. The vegetation in this area is beginning to change to reflect current hydrologic conditions and it is expected that eventually this area will meet the three criteria for a jurisdictional wetland. When included, the total extent of wetland area is approximately 2.22 ha (5.48 ac). This matches very closely with ISGS data [2.2 ha (5.5 ac)] at the 12.5% threshold (Plankell and Pociask 2006; Figure 1).

**Project Goal #2** – A native, non-weedy, emergent wetland community will be created. The performance criterion for this goal states that 1.) at least 50% of the plant species should be non-weedy, native, perennial species and 2.) none of the dominants may be non-native, cattails, or reed canary grass. Neither part of this performance criterion was met in 2006. After the fifth

---

\* In some cases wetland hydrology can be met when a site is inundated or saturated for 5% to 12.5% of the growing season (Environmental Laboratory 1987).

year of monitoring, only 43% (26/60) of the species present were non-weedy, native perennials. Although this value is significantly higher than it was in the first few years, it remains below the 50% threshold.

The dominant species at this created wetland site is *Typha angustifolia*, a perennial, obligate (OBL) wetland species. The non-native narrow-leaf cattail, *Typha angustifolia*, is an aggressive, weedy species. The presence of narrow-leaf cattail was noted in 2002 with an importance value (IV) of 3.3 (Marcum *et al.* 2002). It became one of the dominants in 2003 (IV = 8.73), exploded in 2004 (IV = 41.44) and continued to rise in 2005 (IV = 47.54) (Marcum *et al.* 2002; Marcum and Kurylo 2004; Marcum *et al.* 2005, Marcum *et al.* 2006). In 2006, *Typha angustifolia* was present in 88% of the sample plots and its IV (53.41) increased once again. As mentioned in previous years monitoring reports, *Typha angustifolia* is likely to become even more abundant if management techniques are not employed (Apfelbaum 1985). Management of this area by herbicide or other methods will be necessary if this project goal is to be met. If narrow-leaf cattail is not treated this created wetland site will surely fail to meet Project Goal #2. Weller (1975) and Sale and Wetzel (1983) found cutting cattail prior to flooding achieved good control. Applied Ecological Services and All Services Company (1985) had success with wick and spray applications of Roundup followed by manual clipping of cattail stems. Their treatment was done when cattail seeds were just at the ripening stage.

### Literature Cited

- Apfelbaum, S.I. 1985. Cattail (*Typha* spp.) management. *Natural Areas Journal*. 5(3):9-17.
- Applied Ecological Services and All Services Company. 1985. Report on effects to control cattails (*Typha angustifolia* and *T. latifolia*) at the Swain family pond. Libertyville, Illinois. Unpublished report.
- Brooks, T. 1999. FAP 322 (U.S. 51) mitigation site assessment task order. Memorandum from the Illinois Department of Transportation. 2 pp.
- Brooks, T. 2001. FAP 322 (U.S. 51) monitoring task order. Memorandum from the Illinois Department of Transportation, Springfield. 2p.
- Coopridge, M. and D. Ketzner. 2000. Mitigation Site Assessment for FAP 322 (U.S. 51) in Macon County, Illinois. Technical report submitted to the Illinois Department of Transportation by the Illinois Natural History Survey, Champaign. 20 pp.
- Cox, G.W. 1985. Laboratory manual of general ecology. William C. Brown Press, Dubuque, Iowa.
- Daubenmire, R.F. 1959. A canopy coverage method. *Northwest Science* 33:43-64.
- Doll, J.C. 1990. Soil Survey of Macon County, Illinois. United States Department of Agriculture - Soil Conservation Service in cooperation with the Illinois Agricultural Experiment Station. Soil Report #127.

- Environmental Laboratory. 1987. Corps of Engineers Wetland Delineation Manual. Technical Report Y-87-1, U. S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi. 207 pp.
- Federal Interagency Committee for Wetland Delineations. 1989. Federal Manual for Identifying and Delineating Jurisdictional Wetlands. Cooperative technical publication. U.S. Army Corps of Engineers, U.S. Environmental Protection Agency, U.S. Fish and Wildlife Service, and USDA Soil Conservation Service, Washington, D.C.
- Gleason, H. A. and A. Cronquist. 1991. Manual of the Vascular Plants of the Northeastern United States and Adjacent Canada. The New York Botanical Garden Press, Bronx, New York.
- Grime, J.P. 1974. Vegetation classification by reference to strategies. *Nature*. 250:26-31.
- Grime, J.P., J.G. Hodgson, and R. Hunt. 1988. Comparative plant ecology: a functional approach to common British species. Unwin & Hyman, London.
- Herkert, J.R. and J.E. Ebinger, editors. 2002. Endangered and Threatened Species of Illinois: Status and Distribution. Volume 1 – Plants. Illinois Endangered Species Protection Board, Springfield, Illinois. 161 pp.
- Illinois Department of Transportation. 1996. Wetland Compensation Plan for FAP 322 (US 51) Section 58-20-1 & 58-20-1 (HB, HB-1, HB-2) BR & 48R-1 Macon County, Illinois. 8 pp.
- Marcum, P.B., J. Kurylo, and R. Larimore. 2002. Wetland Mitigation Site Monitoring for FAP 322 (U.S. 51), Macon County, Illinois – 2002. Technical report submitted to the Illinois Department of Transportation by the Illinois Natural History Survey, Champaign, Illinois. 28 pp.
- Marcum, P.B. and J. Kurylo. 2004. Second Year Wetland Mitigation Site Monitoring for FAP 322 (U.S. 51), Macon County, Illinois – 2003. Technical report submitted to the Illinois Department of Transportation by the Illinois Natural History Survey, Champaign, Illinois. 29 pp.
- Marcum, P.B., J. Kurylo, and B. Zercher. 2005. Third Year Wetland Mitigation Site Monitoring for FAP 322 (U.S. 51), Macon County, Illinois – 2004. Technical report submitted to the Illinois Department of Transportation by the Illinois Natural History Survey, Champaign, Illinois. 31 pp.
- Marcum, P.B., J. Kurylo, and B. Zercher. 2006. Fourth Year Wetland Mitigation Site Monitoring for FAP 322 (U.S. 51), Macon County, Illinois – 2005. Technical report submitted to the Illinois Department of Transportation by the Illinois Natural History Survey, Champaign, Illinois. 31 pp.
- Mohlenbrock, R.H. 1986. Guide to the vascular flora of Illinois. Southern Illinois University Press, Carbondale, Illinois. 507 pp.

- Mohlenbrock, R.H. 2002. Vascular flora of Illinois. Southern Illinois University Press, Carbondale, Illinois. 490 pp.
- Plankell, E.T. and G.E. Pociask. 2004. Decatur, U.S. Route 51 Wetland Compensation Site. ISGS #27 in Annual Report for Active IDOT Wetland Compensation and Hydrologic Monitoring Sites. Technical report submitted to the Illinois Department of Transportation by the Illinois State Geological Survey, Champaign, Illinois. pp. 67-73.
- Plankell, E.T. and G.E. Pociask. 2005. Decatur, U.S. Route 51 Wetland Compensation Site. ISGS #27 in Annual Report for Active IDOT Wetland Compensation and Hydrologic Monitoring Sites. Technical report submitted to the Illinois Department of Transportation by the Illinois State Geological Survey, Champaign, Illinois. pp. 29-35.
- Plankell, E.T. and G.E. Pociask. 2006. Decatur, U.S. Route 51 Wetland Compensation Site. ISGS #27 in Annual Report for Active IDOT Wetland Compensation and Hydrologic Monitoring Sites. Technical report submitted to the Illinois Department of Transportation by the Illinois State Geological Survey, Champaign, Illinois. pp.20-26.
- Reed, P.B., Jr. 1988. National list of plant species that occur in wetlands: Illinois. U. S. Fish and Wildlife Service. National Wetlands Inventory. NERC-88/18.13. 117 pp.
- Sale, P.J.M. and R.G. Wetzel. 1983. Growth and metabolism of *Typha* species in relation to cutting treatments. Aquatic Botany. 15:321-334.
- Swink, F.S. and G.S. Wilhelm. 1979. Plants of the Chicago Region. Revised and expanded with keys. 3<sup>rd</sup> edition. Morton Arboretum, Lisle, Illinois.
- Swink, F.S. and G.S. Wilhelm. 1994. Plants of the Chicago region. 4<sup>th</sup> edition. Indiana Academy of Science Press, Indianapolis, Indiana.
- Taft, J.B., G.S. Wilhelm, D.M. Ladd, and L.A. Masters. 1997. Floristic quality assessment for vegetation in Illinois: a method for assessing vegetation integrity. *Ergenia* 15: 3-95.
- Tiner, R.W. 1999. Wetland Indicators: A Guide to Wetland Identification, Delineation, Classification, and Mapping. Lewis Publishers. 392 pp.
- United States Army Corps of Engineers. 1993. Guidelines for developing mitigation proposals. Chicago District. September 1, 1993.
- USDA, NRCS. 2002. The PLANTS Database, Version 3.5 <http://plants.usda.gov>. National Plant Data Center, Baton Rouge, LA 70874-4490 USA.
- Watson, B. and P. Sabatini. 2002. Decatur, U.S. Route 51 Wetland Compensation Site. ISGS #27 in Annual Report for Active IDOT Wetland Compensation and Hydrologic Monitoring Sites. Technical report submitted to the Illinois Department of Transportation by the Illinois State Geological Survey, Champaign, Illinois. pp. 87-93.

Watson, B. and P. Sabatini. 2003. Decatur, U.S. Route 51 Wetland Compensation Site. ISGS #27 in Annual Report for Active IDOT Wetland Compensation and Hydrologic Monitoring Sites. Technical report submitted to the Illinois Department of Transportation by the Illinois State Geological Survey, Champaign, Illinois. pp. 78-84.

Weller, M.W. 1975. Studies of cattail in relation to management of marsh wildlife. Iowa State J. Res. 49(4):383-412.

**Appendix 1.**

**Figure 1. Estimated aerial extent of the created wetland site (FAP 322, U.S. Route 51, Macon County, near Elwin, Illinois). Figure prepared by Brad Zercher, INHS.**

## FAP 322, Mitigation Monitoring Site Macon County




0 400 800 Feet

0 100 200 Meters

 **Wetland site** - 4.96 acres

scale 1:4800  
1 inch=400 ft



01/07

**Appendix 2. Wetland Determination Forms**

## ROUTINE ON-SITE WETLAND DETERMINATION

Site 1 (page 1 of 4)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Marsh

**Legal Description:** NE1/4, NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This marsh is located approximately one mile south of Elwin, IL. It is primarily within the north half of the south infield at the U.S. Route 51-Riley Road interchange.

Do normal environmental conditions exist at this site? Yes: X No:  
Has the vegetation, soils, or hydrology been significantly disturbed? Yes: No: X

### VEGETATION

Dominant Plant Species	Indicator Status	Stratum
1. <i>Typha angustifolia</i>	OBL	herb

*based on quantitative vegetation sampling; Table 3*

Percentage of dominant species that are OBL, FACW, FAC+, or FAC: 100%

**Hydrophytic vegetation:** Yes: X No:

**Rationale:** Greater than 50% of the dominants are OBL, FACW, FAC+, or FAC.

### SOILS

Series and phase: Drummer silty clay loam.

On county hydric soils list? Yes: X No:

Is the soil a histosol? Yes: No: X

Histic epipedon present? Yes: No: X

Redox Concentrations? Yes: X No: Color: 10YR 5/8

Redox Depletions? Yes: X No: Color: 10YR 4/1

Matrix color: 10YR 2/1 over 2.5Y 4/2

Other indicators: Depressional area.

**Hydric soils?** Yes: X No:

**Rationale:** This soil has an iron depleted matrix with iron depletions and concentrations starting below the mollic epipedon and continuing down the profile. Therefore this soil is hydric. This soil also met the F3-Depleted Matrix hydric soil indicator from the NRCS.

## ROUTINE ON-SITE WETLAND DETERMINATION

Site 1 (page 2 of 4)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Marsh

**Legal Description:** NE1/4, NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This marsh is located approximately one mile south of Elwin, IL. It is primarily within the north half of the south infield at the U.S. Route 51-Riley Road interchange.

### HYDROLOGY

Inundated: Yes: X (in part) No: Depth of standing water: up to 0.23 m (9 in)

Depth to saturated soil: saturated to surface over much of the site.

Overview of hydrological flow through the system: This site is located in a depression surrounded by highway embankments. Water enters this site via precipitation, sheetflow from adjacent higher ground (road embankments), and drainage from a culvert under U.S. 51. Water leaves the site primarily via evapotranspiration and slowly through soil infiltration. In addition, water exits the site on the east side and drains into an intermittent stream.

Size of watershed: Approximately 11.17 ha (27.6 ac) (IDOT 1996).

Other field evidence observed: This site has been excavated to hold water for longer periods. Areas of inundation and saturation, bare areas indicating ponded water, algal mats, and mud cracks have been observed at this site.

**Wetland hydrology:** Yes: X No:

**Rationale:** A depressional landscape position and field evidence of saturation and inundation suggest that this site is saturated for a sufficient duration to satisfy the wetland hydrology criterion. This is supported by ISGS hydrology data for this site. Plankell and Pociask (2006) concluded that 2.3 ha (5.7 ac) satisfied the wetland hydrology criterion for greater than 12.5% of the growing season in 2006 (Figure 1).

### DETERMINATION AND RATIONALE:

**Is the site a wetland?** Yes: X No:

**Rationale for decision:** Dominant hydrophytic vegetation, hydric soils, and wetland hydrology are all present; therefore, we determined that this site is a wetland.

Determined by: Paul Marcum (vegetation, hydrology, and GPS)

Jessica Kurylo (soils and hydrology)

Jeff Matthews (vegetation and hydrology)

Brad Zercher (GIS)

Illinois Natural History Survey

Division of Ecology and Conservation Science

1816 S. Oak Street

Champaign, Illinois 61820

(217) 333-8459 (Marcum)

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site 1 (page 3 of 4)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Marsh

**Legal Description:** NE1/4, NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This marsh is located approximately one mile south of Elwin, IL. It is primarily within the north half of the south infield at the U.S. Route 51-Riley Road interchange.

**SPECIES LIST**

Scientific name	Common name	Stratum	Wetland indicator Status	C ♦	Perennial, Annual, Biennial
<i>Abutilon theophrasti</i>	velvet-leaf	herb	FACU-	*	Annual
<i>Acer saccharinum</i>	silver maple	shrub, herb	FACW	1	Perennial
<i>Alisma plantago-aquatica</i>	broad-leaf water-plantain	herb	OBL	2	Perennial
<i>Amaranthus tuberculatus</i>	tall waterhemp	herb	OBL	1	Annual
<i>Ambrosia artemisiifolia</i>	common ragweed	herb	FACU	0	Annual
<i>Asclepias incarnata</i>	swamp milkweed	herb	OBL	4	Perennial
<i>Asclepias verticillata</i>	horsetail milkweed	herb	UPL	1	Perennial
<i>Aster pilosus</i>	hairy aster	herb	FACU+	0	Perennial
<i>Aster puniceus</i>	swamp aster	herb	OBL	7	Perennial
<i>Aster simplex</i>	panicked aster	herb	FACW	3	Perennial
<i>Bidens cernua</i>	nodding beggar's ticks	herb	OBL	2	Annual
<i>Bidens tripartita</i>	beggar's ticks	herb	OBL	2	Annual
<i>Carex vulpinoidea</i>	fox sedge	herb	OBL	3	Perennial
<i>Cassia fasciculata</i>	partridge pea	herb	FACU-	1	Annual
<i>Conyza canadensis</i>	horseweed	herb	FAC-	0	Annual
<i>Cyperus acuminatus</i>	taperleaf flat sedge	herb	OBL	2	Annual
<i>Cyperus esculentus</i>	yellow nut-sedge	herb	FACW	0	Perennial
<i>Cyperus strigosus</i>	straw-colored flatsedge	herb	FACW	0	Perennial
<i>Echinochloa muricata</i>	barnyard grass	herb	OBL	0	Annual
<i>Eleocharis erythropoda</i>	red-rooted spike rush	herb	OBL	3	Perennial
<i>Eupatorium perfoliatum</i>	common boneset	herb	FACW+	4	Perennial
<i>Eupatorium serotinum</i>	late boneset	herb	FAC+	1	Perennial
<i>Euthamia graminifolia</i>	grassleaf goldenrod	herb	FACW-	3	Perennial
<i>Festuca pratensis</i>	meadow fescue	herb	FACU-	*	Perennial
<i>Helenium autumnale</i>	autumn sneezeweed	herb	FACW+	3	Perennial
<i>Hibiscus trionum</i>	flower-of-an-hour	herb	UPL	*	Annual
<i>Hordeum jubatum</i>	squirrel-tail grass	herb	FAC+	*	Perennial
<i>Ipomoea hederacea</i>	ivy-leaved morning glory	herb	FAC	*	Annual
<i>Ipomoea lacunosa</i>	small white morning-glory	herb	FACW	1	Annual
<i>Iris shrevei</i>	southern blue flag	herb	OBL	5	Perennial
<i>Juncus dudleyi</i>	Dudley's rush	herb	FAC	4	Perennial
<i>Juncus tenuis</i>	path rush	herb	FAC	0	Perennial

Species list continued on following page.

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site 1 (page 4 of 4)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Marsh

**Legal Description:** NE1/4, NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This marsh is located approximately one mile south of Elwin, IL. It is primarily within the north half of the south infield at the U.S. Route 51-Riley Road interchange.

**SPECIES LIST**

Scientific name	Common name	Stratum	Wetland indicator Status	C ♦	Perennial, Annual, Biennial
<i>Juncus torreyi</i>	Torrey's rush	herb	FACW	3	Perennial
<i>Leersia oryzoides</i>	rice cutgrass	herb	OBL	3	Perennial
<i>Lobelia cardinalis</i>	cardinal-flower	herb	OBL	6	Perennial
<i>Lobelia siphilitica</i>	blue cardinal-flower	herb	FACW+	4	Perennial
<i>Panicum capillare</i>	witch grass	herb	FAC	0	Annual
<i>Phragmites australis</i>	common red reed	herb	FACW+	1	Perennial
<i>Polygonum pensylvanicum</i>	giant smartweed	herb	FACW+	1	Annual
<i>Polygonum persicaria</i>	spotted lady's thumb	herb	FACW	*	Annual
<i>Populus deltoides</i>	eastern cottonwood	shrub, herb	FAC+	2	Perennial
<i>Potamogeton pectinatus</i>	comb pondweed	herb	OBL	5	Perennial
<i>Pycnanthemum virginianum</i>	common mountain mint	herb	FACW+	5	Perennial
<i>Rumex altissimus</i>	pale dock	herb	FACW-	2	Perennial
<i>Rumex crispus</i>	curly dock	herb	FAC+	*	Perennial
<i>Salix amygdaloides</i>	peach-leaved willow	shrub	FACW	4	Perennial
<i>Salix exigua</i>	sandbar willow	shrub, herb	OBL	1	Perennial
<i>Salix nigra</i>	black willow	tree, shrub	OBL	3	Perennial
<i>Scirpus fluviatilis</i>	river bulrush	herb	OBL	3	Perennial
<i>Scirpus tabernaemontanii</i>	great bulrush	herb	OBL	4	Perennial
<i>Setaria faberi</i>	giant foxtail	herb	FACU+	*	Annual
<i>Setaria glauca</i>	pigeon grass	herb	FAC	*	Annual
<i>Sida spinosa</i>	prickly sida	herb	FACU	*	Annual
<i>Solidago canadensis</i>	Canada goldenrod	herb	FACU	1	Perennial
<i>Solidago gigantea</i>	late goldenrod	herb	FACW	3	Perennial
<i>Solidago riddellii</i>	Riddell's goldenrod	herb	OBL	7	Perennial
<i>Sparganium eurycarpum</i>	burreed	herb	OBL	5	Perennial
<i>Typha angustifolia</i>	narrow-leaved cattail	herb	OBL	*	Perennial
<i>Typha latifolia</i>	cattail	herb	OBL	1	Perennial
<i>Xanthium strumarium</i>	cocklebur	herb	FAC	0	Annual

♦ Coefficient of Conservatism (Taft *et al.* 1997)

\*Non-native species

$$\text{mean C value (mCv)} = \sum C/N = 117/49 = 2.4$$

$$\text{FQI} = \sum C/\sqrt{N} = 117/(\sqrt{49}) = 16.7$$

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site 2 (page 1 of 5)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Non-native Grassland/Forbland

**Legal Description:** NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This non-native grassland/forbland is located approximately 1 mile south of Elwin, IL. It is primarily within the south half of the south infield at the U.S. Route 51-Riley Road interchange. This site also extends around site #1 in the north section of the infield.

Do normal environmental conditions exist at this site? Yes:  No:   
Has the vegetation, soils, or hydrology been significantly disturbed? Yes:  No:

**VEGETATION**

<b>Dominant Plant Species</b>	<b>Indicator Status</b>	<b>Stratum</b>
1. <i>Melilotus officinalis</i>	FACU	herb
2. <i>Solidago canadensis</i>	FACU	herb

*based on quantitative vegetation sampling*

Percentage of dominant species that are OBL, FACW, FAC+, or FAC: 0%

**Hydrophytic vegetation:** Yes:  No:

**Rationale:** Less than 50% of the dominants are OBL, FACW, FAC+, or FAC.

**SOILS**

Series and phase: Flanagan silt loam

On county hydric soils list? Yes:  No:

Is the soil a histosol? Yes:  No:

Histic epipedon present? Yes:  No:

Redox Concentrations? Yes:  No:  Color: 7.5YR 5/8 & 10YR 4/6

Redox Depletions? Yes:  No:

Matrix color: 10YR 2/1 over 10YR 6/6

Other indicators: Higher topographic position relative to wetland site.

**Hydric soils?** Yes:  No:

**Rationale:** A soil matrix color with a chroma greater than 2 is too bright to be considered hydric.

## ROUTINE ON-SITE WETLAND DETERMINATION

Site #2 (page 2 of 5)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Non-native Grassland/Forbland

**Legal Description:** NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This non-native grassland/forbland is located approximately 1 mile south of Elwin, IL. It is primarily within the south half of the south infield at the U.S. Route 51-Riley Road interchange. This site also extends around site #1 in the north section of the infield.

### HYDROLOGY

Inundated: Yes: No: X Depth of standing water: NA

Depth to saturated soil: > 1.2 m (48 in)

Overview of hydrological flow through the system: This site is at a slightly to noticeably higher elevation than site #1. It is level to slightly sloping. Water enters this site via precipitation and sheetflow from adjacent higher ground. Water leaves the site via evapotranspiration, soil infiltration, and through sheetflow into site #1.

Size of watershed: Approximately 11.17 ha (27.6 ac) (IDOT 1996).

Other field evidence observed: none

**Wetland hydrology:** Yes: No: X

**Rationale:** Field observations suggest that this site is both too high of elevation and too sloping to satisfy the wetland hydrology criterion. Furthermore, ISGS hydrology monitoring does not show wetland hydrology for this area (Plankell and Pociask 2006). In our opinion, the site is not saturated long enough during the growing season to meet the wetland hydrology criterion.

### DETERMINATION AND RATIONALE:

<b>Is the site a wetland?</b>	Yes: No: X
<b>Rationale for decision:</b>	Dominant hydrophytic vegetation, hydric soils, and wetland hydrology were all absent; therefore, this site is not a wetland. The National Wetland Inventory did not code this site as a wetland.

Determined by: Paul Marcum and Jeff Matthews (vegetation and hydrology)  
Jessica Kurylo (soils and hydrology)  
Brad Zercher (GPS, GIS and hydrology)  
Illinois Natural History Survey  
Division of Ecology and Conservation Science  
1816 S. Oak Street  
Champaign, Illinois 61820  
(217) 333-8459 (Marcum)

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site #2 (page 3 of 5)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Non-native Grassland/Forbland

**Legal Description:** NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This non-native grassland/forbland is located approximately 1 mile south of Elwin, IL. It is primarily within the south half of the south infield at the U.S. Route 51-Riley Road interchange. This site also extends around site #1 in the north section of the infield.

**SPECIES LIST**

Scientific name	Common name	Stratum	Wetland indicator status	C♦
<i>Abutilon theophrasti</i>	velvet-leaf	herb	FACU-	*
<i>Ambrosia artemisiifolia</i>	common ragweed	herb	FACU	0
<i>Ambrosia trifida</i>	giant ragweed	herb	FAC+	0
<i>Apocynum cannabinum</i>	dogbane	herb	FAC	2
<i>Asclepias incarnata</i>	swamp milkweed	herb	OBL	4
<i>Asclepias syriaca</i>	common milkweed	herb	UPL	0
<i>Asclepias verticillata</i>	horsetail milkweed	herb	UPL	1
<i>Aster pilosus</i>	hairy aster	herb	FACU+	0
<i>Aster puniceus</i>	swamp aster	herb	OBL	7
<i>Aster simplex</i>	panicled aster	herb	FACW	3
<i>Bidens tripartita</i>	beggar's ticks	herb	OBL	2
<i>Bromus inermis</i>	awnless brome grass	herb	UPL	*
<i>Carex normalis</i>	sedge	herb	FACW	4
<i>Cassia fasciculata</i>	partridge pea	herb	FACU-	1
<i>Celtis occidentalis</i>	hackberry	herb	FAC-	3
<i>Chamaesyce nutans</i>	nodding spurge	herb	FACU-	0
<i>Chenopodium album</i>	lamb's quarters	herb	FAC-	*
<i>Cirsium arvense</i>	Canada thistle	herb	FACU	*
<i>Cirsium discolor</i>	pasture thistle	herb	UPL	3
<i>Conyza canadensis</i>	horseweed	herb	FAC-	0
<i>Coreopsis tripteris</i>	tall coreopsis	herb	FAC	4
<i>Cornus drummondii</i>	rough-leaved dogwood	herb	FAC	2
<i>Cornus racemosa</i>	gray dogwood	shrub	FACW-	2
<i>Cynanchum laeve</i>	blue vine	herb	FAC	1
<i>Cyperus esculentus</i>	yellow nut-sedge	herb	FACW	0
<i>Dactylis glomerata</i>	orchard grass	herb	FACU	*
<i>Daucus carota</i>	Queen Anne's lace	herb	UPL	*
<i>Echinacea purpurea</i>	purple coneflower	herb	UPL	6
<i>Elaeagnus umbellata</i>	autumn olive	shrub	UPL	*

Species list continued on following page.

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site #2 (page 4 of 5)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Non-native Grassland/Forbland

**Legal Description:** NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This non-native grassland/forbland is located approximately 1 mile south of Elwin, IL. It is primarily within the south half of the south infield at the U.S. Route 51-Riley Road interchange. This site also extends around site #1 in the north section of the infield.

**SPECIES LIST (continued)**

Scientific name	Common name	Stratum	Wetland indicator status	C♦
<i>Erigeron annuus</i>	annual fleabane	herb	FAC-	1
<i>Eupatorium altissimum</i>	tall boneset	herb	FACU	2
<i>Eupatorium serotinum</i>	late boneset	herb	FAC+	1
<i>Festuca pratensis</i>	meadow fescue	herb	FACU-	*
<i>Geranium carolinianum</i>	wild cranesbill	herb	UPL	2
<i>Helenium autumnale</i>	autumn sneezeweed	herb	FACW+	3
<i>Hibiscus trionum</i>	flower-of-an-hour	herb	UPL	*
<i>Hordeum jubatum</i>	squirrel-tail	herb	FAC+	*
<i>Ipomoea lacunosa</i>	small white morning-glory	herb	FACW	1
<i>Iris shrevei</i>	southern blue flag	herb	OBL	5
<i>Juncus dudleyi</i>	Dudley's rush	herb	FAC	4
<i>Juncus torreyi</i>	Torrey's rush	herb	FACW	3
<i>Juniperus virginiana</i>	eastern red cedar	herb	FACU	1
<i>Lactuca serriola</i>	prickly lettuce	herb	FAC	*
<i>Lobelia cardinalis</i>	cardinal-flower	herb	OBL	6
<i>Lobelia siphilitica</i>	blue cardinal-flower	herb	FACW+	4
<i>Melilotus alba</i>	white sweet clover	herb	FACU	*
<i>Melilotus officinalis</i>	yellow sweet clover	herb	FACU	*
<i>Morus alba</i>	white mulberry	shrub	FAC	*
<i>Oenothera biennis</i>	evening primrose	herb	FACU	1
<i>Oxalis stricta</i>	yellow wood sorrel	herb	FACU	0
<i>Panicum capillare</i>	witch grass	herb	FAC	0
<i>Pastinaca sativa</i>	parsnip	herb	UPL	*
<i>Phragmites australis</i>	common red reed	herb	FACW+	1
<i>Poa pratensis</i>	Kentucky bluegrass	herb	FAC-	*
<i>Polygonum pensylvanicum</i>	giant smartweed	herb	FACW+	1
<i>Polygonum persicaria</i>	spotted lady's thumb	herb	FACW	*
<i>Populus deltoides</i>	eastern cottonwood	shrub	FAC+	2
<i>Pycnanthemum virginianum</i>	common mountain mint	herb	FACW+	5
<i>Ratibida columnifera</i>	long-headed coneflower	herb	UPL	*

Species list continued on following page.

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site #2 (page 4 of 5)

**Field Investigators:** Marcum, Kurylo, and Matthews

**Date:** 5 June and 29 August 2006

**Project Name:** FAP 322 (U.S. 51)

**State:** Illinois

**County:** Macon

**Site Name:** Non-native Grassland/Forbland

**Legal Description:** NE1/4, SE1/4, Section 9, T. 15 N., R. 2 E. and SE1/4, SE1/4, NE1/4, Section 9, T. 15 N., R. 2 E.

**Location:** This non-native grassland/forbland is located approximately 1 mile south of Elwin, IL. It is primarily within the south half of the south infield at the U.S. Route 51-Riley Road interchange. This site also extends around site #1 in the north section of the infield.

**SPECIES LIST (continued)**

Scientific name	Common name	Stratum	Wetland indicator status	C♦
<i>Ratibida pinnata</i>	drooping coneflower	herb	UPL	4
<i>Rudbeckia laciniata</i>	cutleaf coneflower	herb	FACW+	3
<i>Setaria faberi</i>	giant foxtail	herb	FACU+	*
<i>Sida spinosa</i>	prickly sida	herb	FACU	*
<i>Solidago canadensis</i>	Canada goldenrod	herb	FACU	1
<i>Solidago riddellii</i>	Riddell's goldenrod	herb	OBL	7
<i>Taraxacum officinale</i>	common dandelion	herb	FACU	*
<i>Trifolium hybridum</i>	alsike clover	herb	FAC-	*
<i>Trifolium pratense</i>	red clover	herb	FACU+	*
<i>Trifolium repens</i>	white clover	herb	FACU+	*
<i>Verbascum thapsus</i>	woolly mullein	herb	UPL	*

♦ Coefficient of Conservatism (Taft *et al.* 1997)

\*Non-native species

mean C value (mCv) =  $\sum C/N = 113/45 = 2.5$

FQI =  $mCv/(\sqrt{N}) = 113/(\sqrt{45}) = 16.8$

**Appendix 3. Photos of wetland creation sites**



**Photo 1. View from north end of wetland, looking south. A line from the surface water monitoring station (RDS-2) in the foreground to the right side of the road sign in the background (no longer visible because of *Typha angustifolia* expansion) is approximately 205° bearing. This is the location of the baseline established for vegetative sampling (8/29/06).**



**Photo 2. View from east side of the wetland looking toward the *Typha angustifolia* patch near U.S. 51 (8/29/06).**



**Photo 3. Looking south from the middle of the wetland near the north end (8/29/06).**



**Photo 4. View from the south end of the wetland looking north. Note *Festuca pratensis* in the foreground. This is site #2, the non-native grassland (8/29/06).**



**Photo 5. View of site #2, the non-native grassland from near the south end of the wetland (8/29/06).**



**Photo 6. View of *Typha angustifolia* patch looking east from along U.S. 51 (8/29/06).**