

**TRANSMITTAL FORM**

To: Bureau of Design and Environment  
Attn: Thomas Brooks  
From: Illinois Natural History Survey  
Re: Wetland Mitigation Monitoring

**Route and Location**

Mark: La Grange Mitigation Bank Site  
County: Brown  
IDOT District: 6  
Sequence Number: 9579

**Survey Conducted By:** Allen Plocher, Dennis Keene and Brad Zercher  
Division of Ecology and Conservation Science  
Illinois Natural History Survey  
1816 S. Oak St.  
Champaign, IL 61820  
(217) 333-6292

**Date Conducted:** 14, 17 August 2007

**Project Summary:**

We conducted the first year of vegetation monitoring for tree planting in Areas 4 and 7 of the La Grange Mitigation Bank Site. The attached report includes information detailing monitoring methods and results. The status of the created wetland site is discussed. The created wetland site is overlain on digital ortho-quad photography (DOQ) using Arcview 3.2. This report has been posted on the IDOT ftp site as well as submitted as a hard copy.

Signed: \_\_\_\_\_  
Dr. Allen E. Plocher  
INHS/IDOT project Coordinator

Date: \_\_\_\_\_

Signed: \_\_\_\_\_  
Dr. Edward J. Heske  
INHS/IDOT project principal investigator

Date: \_\_\_\_\_

# **Wetland Mitigation Monitoring for the La Grange Mitigation Bank Site, Areas 4 and 7 - 2007**

Allen Plocher, Dennis Keene, Brad Zercher  
Illinois Natural History Survey  
Division of Ecology and Conservation Science  
1816 S. Oak St.  
Champaign, IL 61820  
(217) 333-6292

## **Introduction**

In 2004, the Illinois Department of Transportation (IDOT) established the La Grange Wetland Mitigation Bank in Brown Co., IL (legal location: T. 1 S., R. 1 W., Sect. 16, 17, 20, 21) (Watson et al. 2004). This site, at the confluence of the Illinois and La Moine Rivers, occupies 665 ha (1643 acres), primarily comprising low agricultural fields with some previously existing upland forest, forested wetland, marsh, wet meadow and backwater lakes. Topographically, the site consists of a lower floodplain area, which is inundated for a sufficient duration to support wetland hydrology in more than 7 out of 10 years, a less frequently inundated upper floodplain and a small area of river bluff. The slope break between the lower and upper floodplain occurs at about 132.3 m (434 ft) elevation. To facilitate agriculture, the hydrology of the site has been modified. Ditch and tile drainage systems are in place, a levee is present, and pumps were operational. Since establishment of the bank, the pumps have been removed and portions of the tile and ditch systems deactivated or plugged. In 2002, a flood event breached the levee in two places. For organizational and management purposes, the site has been arbitrarily divided into 16 Areas (Watson et al. 2004).

The general site plan calls for emergent wetland establishment through natural regeneration on the lower floodplain and forested wetland planting on the upper floodplain. Wetland enhancement of areas designated as farmed wetland (FW) is expected to result in 95.8 ha (237 acres) of emergent wetland. Restoration of areas designated as prior converted cropland (PC) is expected to generate 220.3 ha (544.2 acres) of emergent wetland and 117.4 ha (290.1 acres) of forested wetland (Watson et al. 2004). In the fall of 2006, Areas 4 and 7 on the upper floodplain were planted with trees (five foot container grown). Species planted were *Quercus palustris*, *Quercus bicolor*, *Carya illinoensis* and *Carya laciniosa/ovalis*. On 39 acres, 2932 trees were to be planted (actually about 3191 trees were planted), for a proposed stocking of 75/acre. The understory was to be seeded with a mix of *Agrostis alba*, *Phleum pratense*, *Lolium perenne*, *Cinna arundinacea*, *Elymus virginicus*, *Rudbeckia laciniata* and *Polygonum punctatum*. The INHS was tasked to monitor planted tree survival and conduct qualitative assessments of understory vegetation beginning in 2007. This area was in row crops until fall of 2006 (Busemeyer and Plocher 2004).

In 2007, field monitoring was conducted on 14 and 17 August. This report details results

of the 2007 monitoring. Project goals, objectives and performance criteria are included, as are monitoring methods, monitoring results, summary information and recommendations. A wetland banking prospectus (IDOT 2002)) and Wetland Banking Instrument (Watson et al. 2004) were prepared by the Illinois State Geological Survey and Illinois Natural History Survey.

### **Project Goals, Objectives and Performance Criteria**

Proposed goals and objectives are based on information contained in the original IDOT project request (Sunderland, 2006) and the Wetland Banking Instrument (Watson et al. 2004). Performance criteria are based on those specified in the U. S. C. O. E. Wetland Delineation Manual (Environmental Laboratory, 1987), and Guidelines for Developing Mitigation Proposals (USACOE, 1993). Each goal should be attained by the end of the monitoring period. Project goals, objectives and performance criteria are listed below.

**Project goal 1:** The created wetland site should be determined to be jurisdictional by current federal standards.

**Objective:** The goal is to enhance 237 acres of Farmed Wetland and restore 834 acres of Prior Converted cropland by establishing emergent, scrub shrub and forested wetland.

**Performance Criteria:** The entire created wetland should satisfy the three criteria of the federal wetland definition: hydrophytic vegetation, hydric soils and wetland hydrology.

- A. Predominance of hydrophytic vegetation - More than 50% of the dominant plant species must be hydrophytic.
- B. Presence of hydric soils - Hydric soil characteristics must be present, or conditions favorable to the formation of hydric soil must persist at the site.
- C. Presence of wetland hydrology - the created wetland must be inundated at an average depth of less than 2 m (6.6 ft) or have soils saturated to the surface for at least 12.5 % of the growing season.

**Project goal 2:** The created wetland should meet minimum standards as to floristic composition.

**Objective:** The created wetland should compensate in-kind for loss of forested, scrub shrub and emergent wetlands. The wetland compensation should be composed of vegetation characteristic of forested, scrub shrub, and emergent wetlands.

**Performance Criteria:** At least 50% of the planted trees and shrubs should be established and living. At least 75% of the plant species present should be non-weedy, native, annual and perennial species. None of the three most dominant species in any stratum should be nonnative, or weedy species.

## Methods

Monitoring will be performed on the wetland bank site. Illinois Natural History Survey personnel qualitatively monitored the lower floodplain in 2004 and 2005, began quantitative vegetation monitoring in 2006, and will continue until the Illinois Department of Transportation requests that monitoring cease. Monitoring of tree plantings on the upper floodplain began in 2007 and is expected to continue for five years. The Illinois State Geological Survey has been tasked to monitor hydrology. Monitoring reports on the status of the wetland creation site will be submitted annually. The likelihood of meeting the proposed goals and performance criteria will be addressed. If evidence is discovered indicating that the goals/performance criteria will not be met by the end of the monitoring period, written management recommendations will be submitted to IDOT in an effort to correct the problems.

### Project Goal 1

Wetland restoration and enhancement areas will be mapped in the field, and boundaries overlain on digital ortho photographs using Arcview 3.2.

A. Hydrophytic Vegetation - In the lower floodplain area, species composition (relative frequency, relative cover, and Importance Value) will be determined annually through quantitative vegetation sampling of permanent plots. Seventeen parallel transects were established at 200 m (656 ft) intervals. Sampling points (70) were established at 200 m (656 ft) intervals on each transect. At each sampling point, vegetation was tallied by species and percent cover in 20 m<sup>2</sup> quadrats. For Areas 4 and 7 on the upper floodplain, planted trees and natural regeneration will be tallied in 30 m planted row sections at 300 m intervals (10% sample). Beginning when woody vegetation approximates 20% cover, these data will be used to determine woody species composition. Herbaceous species composition in the reforestation areas will be determined using visual estimation. Dominance is based on Importance Value, a numerical average of species' relative frequency, density and/or aerial coverage (Cox 1985). In each stratum dominant species include, starting with the most dominant, those species whose Importance Values, when summed in descending order, exceed 50%, as well as any additional species whose Importance Values are 20% or greater (Federal Interagency Committee for Wetland Delineation, 1989). Dominant species are assigned wetland indicator status ratings (Reed, 1988). Any plant rated facultative or wetter (FAC, FAC+, FACW-, FACW, FACW+ or OBL) is considered hydrophytic. Hydrophytic vegetation is determined to be present if greater than 50% of the dominant species are hydrophytic (Environmental Laboratory 1987).

B. Hydric Soils – In 2000, soil cores collected from the mitigation site were examined for the presence of redoximorphic features (Environmental Laboratory 1987). Being on the floodplain of the Illinois River, virtually the entire area was found to be underlain by hydric soils (IDOT 2002).

C. Wetland Hydrology - The extent of wetland hydrology at this site was monitored by the Illinois State Geological Survey and is shown on the accompanying figure (Fucciolo et al. 2007). Wetland hydrology occurs when inundation or saturation to land surface is present for

greater than 5% of the growing season (10 days at this site) where the soils and vegetation parameters in the Corps of Engineers Wetland Delineation Manual also are met; if either is lacking, then inundation or saturation must be present for greater than 12.5% of the growing season (26 days at this site) to satisfy wetland hydrology criteria (Environmental Laboratory 1987 [<http://el.erdc.usace.army.mil/wetlands/pdfs/wlman87.pdf>]). Inundation and saturation at the site were monitored using a combination of 25 monitoring wells and 8 stage gauges. Water levels were measured at least biweekly during April and May, and monthly during the remainder of the year. Manual readings are generally supplemented by 4 dataloggers, which measure surface and ground-water levels at regular intervals to document all hydrologic events. In 2007, however, no dataloggers were deployed due to extensive on-site flooding, and hence, on-site water level readings were augmented by data from a nearby stream gauging station. Additional details regarding site conditions and monitoring results for wetland hydrology in 2007 are summarized in ISGS' Annual Report for Active IDOT Wetland Compensation and Hydrologic Monitoring Sites, September 1, 2006 to September 1, 2007 (Fucciolo et al. 2007). Information provided by ISGS concerning hydrology of the site is incorporated into this report. In addition, visual inspection of the site for field indicators of wetland hydrology, such as landscape position, inundation or surface saturation or wetland drainage and debris patterns, will be used to determine the presence of wetland hydrology (Environmental Laboratory 1987).

## **Project Goal 2**

A. Tree Survival – Quantitative sampling (30 m planted row sections at 300 m intervals) will be conducted annually to determine whether at least 50% of planted trees remain alive.

B. Vegetation - Dominant plant species in each stratum in each plant community in the lower floodplain area will be determined annually by quantitative sampling. In the upper floodplain area, dominant shrub/sapling layer species will be determined by quantitative sampling, while dominant herbaceous species will be determined by visual estimation. Lists of dominant species will be examined in an attempt to ensure that, in the enhancement and restoration areas, none of the three most dominant species are weedy or non-native. A species list will be prepared annually for each community in order to determine whether at least 75% of the plant species are native and non-weedy. A Floristic Quality Index will be computed annually for each plant community (Taft et al 1997).

## **Results**

**Project Goal 1:** The created wetland site should be determined to be jurisdictional by current federal standards.

In Areas 4 and 7 of the upper floodplain, mowing is carried out several times per year and planted trees are too sparse to constitute a dominant vegetation layer. The herbaceous layer is dominated by *Echinochloa muricata* (OBL) and *Xanthium strumarium* (FAC)

and, therefore, the vegetation is hydrophytic. Areas 4 and 7 of the upper floodplain are underlain by hydric soils (figure 2, Appendix 1).

Although for the past two years this region has experienced relatively severe drought conditions, 1114 acres at the La Grange Bank site (including 23.5 out of 39 acres in Areas 4 and 7), conclusively supported wetland hydrology in 2007 (figure 1). This year these areas were inundated in winter and early spring (Carr et al 2007).

**Former Wessel Property, La Grange Wetland Bank Site**  
**Estimated Areal Extent of 2007 Wetland Hydrology**

map based upon USGS digital orthophotograph, Cooperstown NE quarter quadrangle, produced from 4/14/98 aerial photography (ISGS 2002)




figure 1. Estimated extent of 2007 wetland hydrology

**Project goal 2:** The created wetland should meet minimum standards as to floristic composition.

### Tree Planting

In 2007, winter floods and ice damage uprooted many of the recently planted trees. However, these were fairly quickly replanted and the site did not flood again this year. Therefore tree survival was very good. Area 7 is topographically lower and received more flooding. In this area tree survival was 85.5%. In Area 4 survival was 89.6%. Overall, planted tree survival was 87.5% and, if equal numbers of each species were planted (798), *Quercus palustris* had highest survival (97.3%) and *Carya illinoensis* had the lowest (77.2%). Planted tree survival is well above the required 50% for all species.

### Vegetation

In 2007 this area is only one year out of row crop agriculture and is mowed several times per year. Therefore the vegetation is early successional and weedy. Without planted species, the FQI is 10.3 (14.0 with planted). Although 83% of the species present are native, only 50% are native and nonweedy. The dominant species, *Echinochloa muricata* and *Xanthium strumarium* are both weedy, native species, so the floristic composition goals for this site are not met. On the other hand, none the species present so far are aggressive, persistent weedy species likely to cause management problems. None of the seeded herbaceous species were present in this area in 2007 (Tables 1,2,3,4, Appendix 1, figure 2).

Table 1. Planted Tree Species – Area 4. no. stems, Importance Value (IV), percent survival, n=32

	no. stems	I.V.	percent of 1581
<i>Quercus palustris</i>	389.19	26.94	
<i>Carya illinoensis</i>	379.21	25.90	
<i>Carya laciniosa/ovalis</i>	369.23	24.75	
<i>Quercus bicolor</i>	279.42	22.42	
Total (on 17.8 acres)	1417.05	100.01	89.6%

Table 2. Planted Tree Species – Area 7. no. stems, Importance Value (IV), percent survival, n= 36

	no. stems	I.V.	percent of 1610
<i>Quercus bicolor</i>	462.26	30.86	
<i>Quercus palustris</i>	386.99	27.35	
<i>Carya laciniosa/ovalis</i>	290.24	23.82	
<i>Carya illinoensis</i>	236.45	17.97	
Total (on 21.2 acres)	1375.94	100.00	85.5%

Table 3. Planted Tree Species –Total. no. stems, Importance Value (IV), percent survival, n= 68

	no. stems	I.V.	percent of 3191
<i>Quercus palustris</i>	776.18	27.15	97.3%
<i>Quercus bicolor</i>	741.68	26.43	92.9%
<i>Carya laciniosa/ovalis</i>	659.47	24.37	82.6%
<i>Carya illinoensis</i>	615.66	22.05	77.2%
Total (on 39.0 acres)	2792.99	100.00	87.5%

## Summary and Recommendations

In 2007 this site is only one year out of row crop agriculture. Planted trees have a very good survival rate and far surpass the goal of 50% survival. Out of 39 acres, 23.5 acres conclusively supported wetland hydrology and met the criteria of wetlands. As expected, the mowed groundcover is weedy, with all dominant species weedy natives and only 50% of all species present native and nonweedy. None of the seeded herbaceous species are yet present. However, no problem species are present and the site appears to be doing well at this early stage.

## Literature Cited

- Busemeyer, D. and A. Plocher. 2004. Mitigation Monitoring for La Grange Wetland Bank Site. Report for the IDOT. 3 pp.
- Busemeyer, D. and A. Plocher. 2005. Mitigation Monitoring for La Grange Wetland Bank Site. Report for the IDOT. 6 pp.
- Carr, K., C. Fucciolo, S. Benton, C. Knight, J. Miner, E. Plankell, G. Pociask and B. Sperling. 2007. Annual Report for Active IDOT Wetland Compensation and Hydrologic Monitoring Sites: La Grange Mitigation Bank Site – 2007. Submitted to Illinois Department of Transportation. Illinois State Geological Survey, Champaign, IL. 9 pp.
- Cox, G. W. 1985. Laboratory Manual for General Ecology. 5<sup>th</sup> Edition. W. C. Brown Publishers. Dubuque, IA. 248 pp.
- Environmental Laboratory. 1987. Corps of Engineers Wetland Delineation Manual. Technical Report Y-87-1, U. S. Army Corps of Engineers Waterways Experiment Station, Vicksburg, MI. 207 pp.
- Federal Interagency Committee for Wetland Delineation. 1989. Federal Manual for Identifying and Delineating Jurisdictional Wetlands. USCOE, USEPA, USFWS, USDASCS Cooperative Technical Publication, Washington, DC. 76 pp.
- Illinois Department of Transportation. 2002. Wetland Banking Prospectus, La Grange Site. IDOT, Springfield, IL 156 pp.
- Plocher, A., D. Keene and B. Zercher. 2006. Mitigation monitoring for La Grange wetland bank site. Report for the IDOT. 25 pp.
- Reed, P. B. 1988. National list of plants that occur in wetlands: North Central. USFWS, National Wetlands Inventory. NERC-88/18.13. 117 pp.
- Sunderland, M. 2006. La Grange Wetland Mitigation Bank monitoring task order. Illinois Department of Transportation, Springfield, IL. 3 pp.
- Taft, J., G. Wilhelm, D. Ladd and L. Masters. 1997. Floristic Quality Assessment for vegetation of Illinois: a method for assessing vegetation integrity. *Erignia* 15: 3-95.
- United States Army Corps of Engineers. 1993. Guidelines for developing mitigation proposals. Chicago District.
- Watson, B., D. Busemeyer, K. Carr, A. Plocher and L. Suloway. 2004. Wetland Banking Instrument, La Grange Site. Illinois State Geological Survey and Illinois Natural History Survey, Champaign, IL. 32 pp.

## Appendix 1: Wetland Report

A brief functional assessment of each wetland is provided in this report. However, this assessment is not an exhaustive description of the values of the site. The Floristic Quality Index (FQI), Developed by Taft, Ladd, Wilhelm and Masters (*Floristic Quality Assessment for Vegetation in Illinois*, 1997), was applied to the vegetation of each site. This index should not be used as a substitute for quantitative analysis, but it does provide a measure of floristic integrity. The FQI is calculated as follows:  $I=R/\sqrt{N}$ , where R represents the sum of the numerical ratings for all species recorded in the area, and N represents the number of recorded native species. The mean C is calculated as:  $\text{mean } C=R/N$ . FQI values of 10 or less indicate low natural quality, while sites with values of 20 or more (mean c generally greater than 3.0) have at least some evidence of native character and may be considered environmental assets.

---

Site 1: This wet meadow is located in the mowed, tree planting of Areas 4 and 7. Hydrophytic vegetation, hydric soils and wetland hydrology are all present. Therefore this site is a wetland. The site occupies 15.8 ha (39 acres), of which 9.5 ha (23.5 acres) conclusively supported wetland hydrology in 2007. Hydrologic inputs are precipitation, sheetflow and overflow from the Illinois River. Water leaves the site by evapotranspiration and sheetflow. The site provides floodwater storage and wildlife habitat of fair quality. The NWI does not code the site as wetland. The FQI is 14.0, which is indicative of fair natural quality.

### Watershed Data:

This site is in the watershed for the Illinois River, which has a drainage area of 62,748 km<sup>2</sup> (24,227 mi<sup>2</sup>) at Beardstown, IL. The USGS hydrologic unit code is 07130011, Illinois River, Lower.

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site 1 (page 1 of 4)

**Field Investigators:** Plocher, Keene, Zercher      **Date:** 14, 17 August 2007  
**Project Name:** LaGrange/Brown Co. Mitigation Bank  
**State:** Illinois    **County:** Brown      **Applicant:** IDOT District 6  
**Site Name:** Wet Meadow  
**Legal Description:** T. 1 S., R. 1 W., E/2 Sect. 17

**Location:** Areas 4 and 7

Do normal environmental conditions exist at this site?      Yes: X    No:  
Has the vegetation, soil, or hydrology been significantly disturbed?    Yes:      No: X

**VEGETATION**

<b>Dominant Plant Species</b>	<b>Stratum</b>	<b>Indicator Status</b>
1. <i>Echinochloa muricata</i>	herb	OBL
2. <i>Xanthium strumarium</i>	herb	FAC

Percent of dominant species that are OBL, FACW, FAC+, or FAC: 100%

**Hydrophytic vegetation:** Yes: X    No:

**Rationale:** More than 50% of dominants are OBL, FACW, FAC+, or FAC.

**SOILS\***

\* field checked in 2000

Series and phase: Mapped as Beaucoup silty clay loam, Titus silty clay loam and Wagner silt loam by NRCS. Revised to Wagner silt loam

On county hydric soils list?      Yes: X    No:  
Is the soil a histosol?              Yes:      No: X  
Histic epipedon present?          Yes:      No: X  
Redox Concentrations?              Yes: X    No:  
Redox Depletions?                  Yes: X    No:

Matrix color: N 4/ and 5Y 5/1

Other indicators: level landscape position

**Hydric soils?**      Yes: X    No:

**Rationale:** This soil meets the requirements for NRCS hydric soil indicators F2 – loamy gleyed matrix, F3 – depleted matrix.

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site 1 (page 2 of 4)

**Field Investigators:** Plocher, Keene, Zercher      **Date:** 14, 17 August 2007  
**Project Name:** LaGrange/Brown Co. Mitigation Bank  
**State:** Illinois    **County:** Brown      **Applicant:** IDOT District 6  
**Site Name:** Wet Meadow  
**Legal Description:** T. 1 S., R. 1 W., E/2 Sect. 17

**Location:** Areas 4 and 7

**HYDROLOGY**

Inundated: Yes:    No: X      Depth of standing water: NA

Depth to saturated soil: > 0.66 m (26 in)

Overview of hydrological flow through the system: Primary hydrologic inputs to this site are precipitation, sheetflow and overflow from the Illinois River. Evapotranspiration and sheetflow are the major outputs.

Size of watershed: 62,748 km<sup>2</sup> (24,227 mi<sup>2</sup>) at Beardstown, IL

Other field evidence observed: level landscape position

**Wetland hydrology:** Yes: X No:

**Rationale:** Field evidence listed above indicates that this site is flooded or saturated for a sufficient period during the growing season to meet the criterion of wetland hydrology.

**WETLAND DETERMINATION AND RATIONALE:**

**Is the site a wetland?:** Yes: X      No:

**Rationale:** Hydrophytic vegetation, hydric soils and wetland hydrology are all present. Therefore the site is a wetland. The site is not coded as wetland by the NWI.

Determined by: Allen Plocher (vegetation and hydrology)  
Dennis Keene (soils and hydrology)  
Brad Zercher (GPS and hydrology)  
Illinois Natural History Survey  
Division of Ecology and Conservation Science  
1816 S. Oak St.  
Champaign, Illinois 61820  
(217) 333-6292

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site 1 (page 3 of 4)

**Field Investigators:** Plocher, Keene, Zercher      **Date:** 14, 17 August 2007  
**Project Name:** LaGrange/Brown Co. Mitigation Bank  
**State:** Illinois    **County:** Brown      **Applicant:** IDOT District 6  
**Site Name:** Wet Meadow  
**Legal Description:** T. 1 S., R. 1 W., E/2 Sect. 17

**Location:** Areas 4 and 7

SPECIES LIST

Scientific name	Common name	Stratum	Wetland indicator status	C*
<i>Abutilon theophrasti</i>	velvet leaf	herb	FACU-	
<i>Acer saccharinum</i>	silver maple	seedling	FACW	1
<i>Amaranthus tuberculatus</i>	water hemp	herb	OBL	1
<i>Ambrosia trifida</i>	giant ragweed	herb	FAC+	0
<i>Ammannia coccinea</i>	ammannia	herb	OBL	5
<i>Apocynum cannabinum</i>	dogbane	herb	FAC	2
<i>Artemisia annua</i>	annual wormwood	herb	FACU	
<i>Asclepias syriaca</i>	common milkweed	herb	UPL	0
<i>Aster simplex</i>	panicled aster	herb	FACW	3
<i>Bidens aristosa</i>	swamp marigold	herb	FACW	1
<i>Bidens connata</i>	beggar's ticks	herb	OBL	2
<i>Bidens frondosa</i>	beggar's ticks	herb	FACW	1
<i>Boltonia asteroides</i>	false aster	herb	FACW	5
<i>Calystegia sepium</i>	hedge bindweed	herb	FAC	1
<i>Campsis radicans</i>	trumpet creeper	herb	FAC	2
<i>Carex lacustris</i>	lake sedge	herb	OBL	6
<i>Carya illinoensis</i>	pecan	shrub	planted	6
<i>Carya laciniosa</i>	shell bark hickory	shrub	planted	7
<i>Carya ovalis</i>	sweet pignut hickory	shrub	planted	5
<i>Cassia fasciculata</i>	partridge pea	herb	FACU-	1
<i>Cephalanthus occidentalis</i>	buttonbush	shrub	OBL	4
<i>Chamaesyce maculata</i>	nodding spurge	herb	FACU-	0
<i>Chenopodium album</i>	lamb's quarters	herb	FAC-	
<i>Conyza canadensis</i>	horseweed	herb	FAC-	0
<i>Cyperus ferruginescens</i>	flat sedge	herb	OBL	1
<i>Digitaria ischaemum</i>	smooth crabgrass	herb	FACU	
<i>Echinochloa muricata</i>	barnyard grass	herb	OBL	0

\*Coefficient of Conservatism, as developed by Taft, Ladd, Wilhelm and Masters (1997)

Continued on following page

**ROUTINE ON-SITE WETLAND DETERMINATION**

Site 1 (page 4 of 4)

**Field Investigators:** Plocher, Keene, Zercher      **Date:** 14, 17 August 2007  
**Project Name:** LaGrange/Brown Co. Mitigation Bank  
**State:** Illinois    **County:** Brown      **Applicant:** IDOT District 6  
**Site Name:** Wet Meadow  
**Legal Description:** T. 1 S., R. 1 W., E/2 Sect. 17

**Location:** Areas 4 and 7

SPECIES LIST (Continued)

Scientific name	Common name	Stratum	Wetland indicator status	C*
<i>Eleocharis obtusa</i>	spikerush	herb	OBL	2
<i>Eragrostis pectinacea</i>	Carolina lovegrass	herb	FAC	0
<i>Erechtites hieracifolia</i>	fireweed	herb	FACU	2
<i>Erigeron annuus</i>	annual fleabane	herb	FAC-	1
<i>Eupatorium serotinum</i>	late flowering thoroughwort	herb	FAC+	1
<i>Fragaria virginiana</i>	wild strawberry	herb	FAC-	2
<i>Ipomoea hederacea</i>	ivy leaf morning glory	herb	FAC	
<i>Ipomoea lacunosa</i>	small flowered morning glory	herb	FACW	1
<i>Lysimachia ciliata</i>	fringed loosestrife	herb	FACW	4
<i>Oenothera biennis</i>	evening primrose	herb	FACU	1
<i>Panicum capillare</i>	witchgrass	herb	FAC	0
<i>Panicum dichotomiflorum</i>	fall panicum	herb	FACW-	0
<i>Phytolacca americana</i>	pokeweed	herb	FAC-	1
<i>Plantago rugelii</i>	Rugel's plantain	herb	FAC	0
<i>Polygonum amphibium</i>	water smartweed	herb	OBL	3
<i>Polygonum lapathifolium</i>	nodding smartweed	herb	FACW+	0
<i>Polygonum pensylvanicum</i>	giant smartweed	herb	FACW+	1
<i>Potentilla norvegica</i>	rough cinquefoil	herb	FAC	0
<i>Quercus bicolor</i>	swamp white oak	shrub	planted	7
<i>Quercus palustris</i>	pin oak	shrub	planted	4
<i>Rumex crispus</i>	curly dock	herb	FAC+	
<i>Setaria faberi</i>	giant foxtail	herb	FACU+	
<i>Sida spinosa</i>	prickly sida	herb	FACU	
<i>Stachys tenuifolia</i>	slenderleaf betony	herb	OBL	5
<i>Trifolium pratense</i>	red clover	herb	FACU+	
<i>Ulmus americana</i>	American elm	seedling	FACW-	5
<i>Xanthium strumarium</i>	cocklebur	herb	FAC	0

\*Coefficient of Conservatism, as developed by Taft, Ladd, Wilhelm and Masters (1997)

FQI (with planted species) =  $R/\sqrt{N} = 94/\sqrt{45} = 14.0$ , mean C =  $R/N = 94/45 = 2.09$

FQI =  $R/\sqrt{N} = 65/\sqrt{40} = 10.3$ , mean C =  $R/N = 65/40 = 1.63$

Percent native species = 83.3

Percent nonweedy and native = 50.0

# LaGrange Mitigation Site, Areas 4 and 7 Brown County



 Site boundary

0 400 Feet

0 100 Meters

scale 1:4800  
1 inch=400 ft

